# THATION PAGE

Form Approved OMB No. 0704-018

AD-A277 003 

Nashington Headquarters Services, Directorate for into Paperwork Reduction Project (0/04-0188) Washington UC 20503

ORT DATE March 1994 3. REPORT TYPE AND DATES COVERED

Interim, 1 July 1992 - 1 March 1994

4. TITLE AND SUBTITLE

Tetraalkyldiarsines as Potential Precursors for Electronic Materials: Synthesis and Characterization of Various Iso-Propyl Arsenic Compounds 5. FUNDING NUMBERS

G: N00014-92-J-1828 R&T PR: 44135035---01

Lawrence F. Brough, Liu Gang, Matthew A. Lipkovich, Thomas J. Colacot, Virgil L. Goedken, and William S. Rees, Jr.

> PERFORMING ORGANIZATION REPORT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

School of Chemistry and Biochemistry and School of Materials Science and Engineering Georgia Institute of Technology Atlanta. Georgia 30332-0400

TR No. 4

9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS

Department of the Navy Office of the Chief of Naval Research Arlington, Virginia 22217-500

10. SPONSORING/MONITORING AGENCY REPORT NUMBER

unknown

11. SUPPLEMENTARY NOTES

Accepted for Publication in: Materials Research Society Proceedings.

12a. DISTRIBUTION / AVAILABILITY STATEMENT

This document has been approved for public release and sale; its distribution is unlimited.

94-08229

13. ABSTRACT (Maximum 200 words)

Tetrakis(iso-propyl)diarsine was synthesized by the reaction of (i-Pr)AsLi with (i-Pr)AsI. The lithium salt of the secondary arsine was produced following deprotonation of (i-Pr)<sub>2</sub>AsH, obtained by reduction of  $(i-Pr)_2$ AsI, which was prepared by the thermolysis of  $(i-Pr)_2$ AsI<sub>2</sub>. The X-ray crystal structure of [(i-Pr)<sub>3</sub>AsI][I] has been determined on the product of the reaction of  $(i-Pr)_3$ As and  $I_2$ . Compounds of the general form  $E=As(i-Pr)_3$  (E = O, S, Se) have been prepared.

رلج فنقداله سراء والمساور

organoarsenic chemistry, iso-propyl arsenic compounds, diarsines			15. NUMBER OF PAGES
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	UL

### OFFICE OF NAVAL RESEARCH

GRANT N00014-92-J-1820

R&T Code 4135035---01

Technical Report No. 4

Tetraalkyldiarsines as Potential Precursors for Electronic Materials: Synthesis and Characterization of Various Iso-Propyl Arsenic Compounds

by

Lawrence F. Brough, Liu Gang, Matthew A. Lipkovich, Thomas J. Colacot, Virgil L. Goedken, and William S. Rees, Jr.

Accepted for Publication in

Materials Research Society Proceedings

School of Chemistry and Biochemistry and School of Materials Science and Engineering Georgia Institute of Technology Atlanta, Georgia 30332-0400

8 March 1994

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited.

Accession For

HTIS GRANI
DTIC TAB
Unamounced
JESTIFICATION

By
DESCRIPTION

Availability Coles

Availability Coles

Dist Special

Tetraalkyldiarsines as Precursors in the Synthesis of Electronic Materials: Synthesis and Characterization of Various iso-Propyl Arsenic Compounds

Lawrence F. Brough, Liu Gang, Matthew A. Lipkovich.

Thomas J. Colacot, Virgil L. Goedken, and William S. Rees, Jr.\*

Department of Chemistry

and Materials Research and Technology Center

The Florida State University

Tallahassee, Florida 32306-3006

VOICE: 904-644-4768

FAX: 904-644-8281

Fall Meeting of the Materials Research Society
Boston, MA

Symposium W

Manuscript Number: W 2.8

# TETRAALKYLDIARSINES AS POTENTIAL PRECURSORS FOR ELECTRONIC MATERIALS: SYNTHESIS AND CHARACTERIZATION OF VARIOUS ISO-PROPYL ARSENIC COMPOUNDS

LAWRENCE F. BROUGH, LIU GANG, MATTHEW A. LIPKOVICH, THOMAS J. COLACOT, VIRGIL L. GOEDKEN,‡ AND WILLIAM S. REES, JR.

Department of Chemistry and Materials Research and Technology Center. The Florida State University, Tallahassee, Florida 32306-3006

#### **ABSTRACT**

Tetrakis(iso-propyl)diarsine was synthesized by the reaction of  $(i-Pr)_2$ AsLi with  $(i-Pr)_2$ AsI. The lithium salt of the secondary arsine was produced following deprotonation of  $(i-Pr)_2$ AsI, obtained by reduction of  $(i-Pr)_2$ AsI, which was prepared by the thermolysis of  $(i-Pr)_3$ AsI<sub>2</sub>. The X-ray crystal structure of  $[(i-Pr)_3$ AsI][I] has been determined on the product of the reaction of  $(i-Pr)_3$ As and I<sub>2</sub>. Compounds of the general form E=As $(i-Pr)_3$  (E = O, S, Se) have been prepared.

### INTRODUCTION

Several recent publications have discussed the potential of utilizing tetrakis(alkyl)diarsines as in situ sources of arsenic for employment in the preparation of arsenic-containing electronic materials. Such routes to, for example, GaAs presently rely on toxic and highly volatile precursors (e.g., AsH<sub>3</sub>). The motivation for this new approach is depicted in equation 1. Although the molecularity of the arsenic-containing product is uncertain, such routes have seen some moderate success. The precise stoichiometry of the reaction product may be 1/2 (As<sub>4</sub>) or 2 As; however, the confirmation of this synthetic procedure for in situ generation of elemental arsenic-containing species relies less on the accretion coefficient, and more on the lack of any carbon-containing species in the product. Such contamination by the disadvantageous decomposition of pendant organic groups highly is detrimental to the final electronic properties of such materials.

<sup>&</sup>lt;sup>‡</sup>Deceased 22 December 1992


<sup>\*</sup>Address all correspondence to this author at: School of Chemistry and Biochemistry and School of Materials Science and Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332.

As one component of a program directed at an examination of the generality of this approach, lc several *iso*-propyl arsenic compounds have been prepared. The synthesis and characterization of some of these organometallic compounds is reported in this contribution.

$$3 R_2 As - As R_2 \qquad \qquad 4 As R_3 + As_2 \tag{1}$$

# RESULTS AND DISCUSSION

Credit is given to Cadet for the initial report of a compound containing a metal-carbon bond.<sup>2</sup> His 1760 paper describes the simplest tetraalkyldiarsine,  $(H_3C)_2As-As(CH_3)_2$ , given the trivial name of cacodyl. A variety of methods have been developed over the years for the preparation of diarsines.<sup>3</sup> In order to address the above-described motivation for exploration of tetraalkyldiarsines as potential precursors in the OMVPE growth of high-quality semiconducting materials, tetrakis(iso-propyl)diarsine was prepared (Scheme I).



Scheme I

# Crystal structures of [(i-Pr)3AsI][I] and [(i-Pr)3AsI] [I3]

The general format for the oxidative addition of a dihalogen molecule to a tertiary organoarsine (equation 2) can proceed according to two alternate pathways.

Route A 
$$R_3AsX_2$$

Route B  $[R_3AsX][X]$ 

(2)

In Route A, the  ${}^{\sim}T_d$  R<sub>3</sub>As is transformed into the molecular R<sub>3</sub>AsX<sub>2</sub> with a trigonal bipyramidal geometry. In Route B, the resultant product is an [arsonium][halide] ionic moiety. Factors influencing the path of choice include the electronic and spatial considerations of the alkyl groups represented in the tertiary arsine, as well as the  $\Delta\chi$  present between the halogen and arsenic. For example, both [Me<sub>3</sub>PX][X] and Me<sub>3</sub>SbX<sub>2</sub> have been characterized.<sup>4</sup> In all likelyhood, the difluorides are covalently bound. Previous infrared spectroscopic work has indicated that R<sub>3</sub>AsI<sub>2</sub> compounds (R = Me, Et) are ionic in the solid state.<sup>5</sup> In the present investigation, [(*i*-Pr)<sub>3</sub>AsI][I] was determined, by X-ray diffraction, to be the conformation adopted by the reaction product of (*i*-Pr)<sub>3</sub>As and I<sub>2</sub>. If the stoichiometry was not carefully controlled (equation 3), a different product was obtained. There have been reports of comparable reactions observed for Ph<sub>3</sub>As.<sup>6</sup>

$$[(i-Pr)_3AsI][I] + I_2 \qquad \qquad [(i-Pr)_3AsI][I_3]$$
 (3)

A single crystal structure determination also was conducted on [(*i*-Pr)<sub>3</sub>AsI][I<sub>3</sub>]. It was, however, suspended prior to complete refinement. Once the identical atomic linkages were ascertained to be present in the organometallic cation, the structural refinement was not pursued further. There was little new structural information revealed in the anion replacement species.

Pertinent crystal and refinement data along with selected interatomic distances and angles will be published elsewhere. An ORTEP representation of the structure is presented (Figure 1), including both covalent and ionic interactions between As and I atoms.




Figure 1 ORTEP representation of [(*i*-Pr)3AsI][I] Open circles represent Carbon atoms. Hydrogen atoms have been omitted for clarity of presentation.

# Preparation of $E=As(i-Pr)_3$ (E = O, S, Se)

The preparation of  $O=As(i-Pr)_3$  was effected under mild conditions (equation 4).

$$As(i-Pr)_3 + \frac{1}{2}H_2O_2 \longrightarrow O=As(i-Pr)_3$$
 (4)

Either a solvent-mediated or a direct reaction can be utilized in the synthesis of S=As(i-Pr)<sub>3</sub> (equation 5 - 6).

$$^{1}/_{8}S_{8} + As(i-Pr)_{3} = \frac{THF, r.t.}{N_{2(g)}, 18 \text{ h}} S=As(i-Pr)_{3}$$
 (5)

$$^{1}/_{8}S_{8} + As(i-Pr)_{3} \xrightarrow{\text{neat}} S=As(i-Pr)_{3}$$
 (6)

The most efficient route for the preparation of  $Se=As(i-Pr)_3$  was determined to be the reaction of  $As(i-Pr)_3$  with elemental selenium (equation 7).

$$As(i-Pr)_3 + \frac{1}{x} Se_x = \frac{150^{\circ}C}{10 \text{ min}}$$
 Se= $As(i-Pr)_3$  (7)

# **ACKNOWLEDGEMENTS**

We acknowledge support of this project by the Office of Naval Research Chemistry Division. L.F.B. was the recipient of a Florida State University Summer Faculty Professional Enhancement Program stipend during the summer of 1990.

# REFERENCES

- a) C. W. Porter and P. Borgstrom, J. Am. Chem. Soc. 41, 2048 (1919). b) Wide Gap II-VI Semiconductors, W. Kuhn, H. P. Wagner, H. Stanzi, K. Wolf, K. Worle, S. Lankes, J. Betz, M. Woraz, D. Lichtenberger, H. Leiderer, W. Gebhard and R. Triboulet, R. L. Aulombard and J. B. Mullin, eds., (Adam Hilger, Bristol, UK), 1991. p. A105. c) V. L. Goedken, L. F. Brough and W. S. Rees, Jr., J. Organomet. Chem. 449, 125 (1993).
- 2. C. Elschenbroich and A. Salzer, <u>Organometallics: A Concise Introduction</u>, (VCH, New York), 1992.
- 3. G. O. Doak and L. D. Freedman, <u>Organometallic Compounds of Arsenic, Antimony, and Bismuth</u>, (Wiley-Interscience, New York), 1970.
- 4. a) J. Goubeau and R. Baumgärtner, Z. Electrochem. **64**, 598 (1960). b) E. F. Wells, Z. Kristallogr. **99**, 367 (1938).
- 5. a) M. H. O'Brien, G. O. Doak, and G. G. Long, Inorg. Chim. Acta 1, 34 (1967). b) M. H. O'Brien, Ph.D. Thesis, North Carolina State University, 1968.
- 6. a) K. R. Bhaskar, S. N. Bhat, S. Singh, and C. N. R. Rao, J. Inorg. Nucl. Chem. 28, 1915 (1966). b) S. N. Bhat and C. N. R. Rao, J. Amer. Chem. Soc. 88, 3216 (1966).

# TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)\*
Chemistry Division, Code 1113
800 North Quincy Street
Arlington, Virginia 22217-5000

Dr. James S. Murday (1) Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)
Chemistry Division, Code 385
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Elek Lindner (1)
Naval Command, Control and Ocean
Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1) Crane Division Naval Surface Warfare Center Crane, Indiana 47522-5000 Dr. Richard W. Drisko (1)
Naval Civil Engineering
Laboratory
Code L52
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Defense Technical Information Center (2) Building 5, Cameron Station Alexandria, VA 22314

\* Number of copies to forward